

RESEARCH ARTICLE

Does Etiology of Acute Pancreatitis Matter? A Review of 391 Consecutive Episodes

Gunther Weitz, Julia Woitalla, Peter Wellhöner, Klaus Schmidt, Jürgen Büning, Klaus Fellermann

Gastroenterology, Medical Department I, University Hospital of Schleswig-Holstein, Campus Lübeck.
Lübeck, Germany

ABSTRACT

Context Acute pancreatitis can be triggered by a variety of factors ranging from short lasting to sustained disruptions. It is plausible that the characteristics and course of disease differ among etiologies. Data distinguishing characteristics of patients with pancreatitis of biliary, alcoholic, idiopathic or other origin are scarce and conflicting. **Objective** To compare patients' characteristics, baseline parameters on admission, and outcome in patients with an episode of acute pancreatitis in whom the etiology was thoroughly determined. **Design** Retrospective study. **Setting** Single center. **Patients** Three-hundreds and 91 consecutive episodes of acute pancreatitis through the years 2008 to 2011. **Main outcome measures** Gender, age, body mass index, Charlson comorbidity index, history of pancreatitis, heart rate, blood pressure, plasma lipase, hematocrit, plasma creatinine, white blood cell count, rate of persistent organ failure and necrosis, maximum C-reactive protein, duration of hospitalization, mortality. **Results** There were marked differences between the groups. Biliary etiology was associated with higher age and body weight, female predominance, higher plasma lipase, and a favourable outcome. Alcoholic etiology had male predominance, a tendency for initial hemoconcentration, a lower plasma lipase, and the highest rate of necrosis. Idiopathic etiology had the highest rate of persistent organ failure and the highest mortality. **Conclusions** Biliary, alcoholic and idiopathic acute pancreatitis should be treated as distinct entities. While alcoholic episodes have the highest risk of necrosis, the worst outcome was observed in the idiopathic group. Hence, finding no causality for an episode of acute pancreatitis after thorough investigation might be a predictor for poor outcome. Larger studies are warranted to confirm this.

INTRODUCTION

Acute pancreatitis can be triggered by a variety of factors including duct obstruction, toxic agents, primary inflammation (e.g. infection, autoimmune), and trauma [1, 2]. The most common etiologies are gallstones and alcohol comprising 60-85% of all incidents [3-5]. Approximately 10-20% occurs without a detectable cause and are considered "idiopathic". However, the pathophysiological pathway from trigger to the destructive response has not been entirely determined yet. It is currently not known whether etiology has an impact on the course of disease. However, differences may be relevant in terms of severity prediction, monitoring, and treatment of patients with acute pancreatitis.

Some differences between the etiologic groups are well recognized: The incidence of acute biliary pancreatitis increases with age and has a female predominance.

Patients with alcoholic pancreatitis are younger and by far more often male [6, 7]. However, published data on the impact of etiology on outcome in acute pancreatitis are conflicting. While some authors deny a relationship others describe biliary pancreatitis to be either most or less severe [8-11]. One study observed patients with alcohol-induced pancreatitis and initial organ failure at highest risk for further deterioration [12]. Another publication revealed high rates of persistent organ failure in alcoholic pancreatitis [13]. In a population-based study, patients with alcoholic pancreatitis had the highest risk for a fatal outcome [7]. When the model was adjusted for age, race and gender the mortality risk was highest for idiopathic pancreatitis. A high mortality in idiopathic has been reported in other studies [14-16].

Several issues limit the comparability of studies on etiology of acute pancreatitis. Firstly, the etiology is often difficult to determine. In patients with presumed idiopathic acute pancreatitis, endoscopic ultrasound (EUS) and magnetic-resonance cholangiopancreatography (MRCP) upon follow-up found possible causes in the majority of cases (mostly cholelithiasis and biliary sludge) [17]. Secondly, there are vast differences in incidence of the etiologies in different countries. E.g., while 24% biliary and 61% alcoholic cases are reported in Hungary, the respective proportions are 71% and 6% in Greece and 60% and 13% in Italy [3]. Thirdly, both severities on admission and case fatality have reduced during the last decades [4]. Moreover, recent advances in the treatment of infected necrosis

Received January 10th, 2015 – **Accepted** February 28th, 2015

Keywords Pancreatitis /etiology; Retrospective Studies

Abbreviations CCI: Charlson comorbidity index; CRP: C-reactive protein; ERC: endoscopic retrograde cholangiography; EUS: endoscopic ultrasound

Correspondence Gunther Weitz

University Hospital of Schleswig-Holstein, Campus Lübeck
Medical Department I, Gastroenterology
Ratzeburger Allee 160, D-23538 Lübeck, Germany

Phone ++49 4515006033

Fax ++49 4515006242

E-mail gunther.weitz@uksh.de

might have helped to further reduce mortality [18]. Hence, the actual characteristics of the different etiological types of acute pancreatitis are uncertain.

We therefore sought to analyze the characteristics of the three most frequent types of acute pancreatitis in consecutive episodes of the disease with thoroughly determined etiology. Our hypothesis was that we could detect differences in the characteristics that could be relevant for prognosis and therapy.

METHODS

We reviewed the charts of all our inpatients, who presented with acute pancreatitis in our emergency department from January 2008 to December 2011. All patients were at least 18-year-old. Acute pancreatitis was defined by the revised Atlanta Classification [19]. Referrals from other hospitals were excluded. The tracking process excluding patients with false disease identification and missing or incomplete charts and the standard treatment in our department have been described elsewhere in detail [20].

Biliary etiology was defined as evidence of a gallstone in the bile duct and/or a cholestatic pattern of liver function tests (alanine and aspartate transaminase, gamma-glutamyltransferase, alkaline phosphatase) normalizing within days after admission. All patients underwent abdominal ultrasound testing. In patients with cholangitis or worsening jaundice, a prompt endoscopic retrograde cholangiography (ERC) with sphincterotomy was performed. This procedure was also electively performed in patients that were too old or too ill for cholecystectomy. All other patients underwent EUS a few days after the event to evaluate the bile duct and the pancreas. In case of retained gall stones in the common bile duct, an ERC with sphincterotomy was performed. An early cholecystectomy was recommended in all eligible patients with biliary pancreatitis. Commonly, patients were discharged and free to choose a hospital for surgery. Data on the cholecystectomy rate and the date of surgery were not available. Alcoholic etiology was defined as evidence of sustained uncontrolled consumption of alcoholic beverages (>30 g alcohol per day in men and >20 g alcohol per day in women) and/or an alcohol excess (defined as a drunken state in the patient's history) in the week prior to admission while excluding acute pancreatitis of other causes.

All patients were screened for hypercalcemia, hyperlipidemia, hereditary conditions, and in unclear cases for Immunoglobulin G4 antibodies, lactoferrin antibodies, anticarbonic anhydrase antibodies, and anti-nuclear antibodies. Moreover, all patients that were primarily considered non-biliary underwent endoscopic ultrasound (EUS) a few days after the event to evaluate the pancreas and the bile duct. Thereby they were screened for local complications and for etiological hints such as tumors or pathognomonic pancreas morphology. In case of incidentally found calculi in the bile duct, the acute pancreatitis was considered biliary and was treated as

mentioned above. Chronic pancreatitis was diagnosed when imaging showed evidence of the disease. Patients with evidence of necrosis or who did not recover within four days after admission underwent a contrast enhanced CT scan. Local and systemic complications were defined by the revised Atlanta Classification [19].

ETHICS

Data was extracted from the charts and further processed without the identification of the patients. The study was approved by means of the local ethics committee. The work was carried out in accordance with the Declaration of Helsinki, and the anonymity of all participants was guaranteed.

STATISTICS

The presentation of the data is descriptive. The differences between continuous variables of the four etiology groups were tested by the Kruskal-Wallis test, the categorical variables were tested by Pearson's chi-squared test. A P value less than 0.05 was considered significant.

RESULTS

We identified 391 episodes of acute pancreatitis, 145 (37.1%) biliary, 123 (31.5%) alcoholic, 61 (15.6%) idiopathic and 62 (15.9%) others. One hundred and twenty-two episodes (31.2%) in 78 patients were recurrent. In 68 cases (17.4%) there was evidence of chronic pancreatitis. In 54 of the 391 episodes (13.8%) a necrosis was detected. Nine patients (6 alcoholic, 3 idiopathic) were treated with drainage (8 transgastral, 1 percutaneous). One of the patients with transgastral drainage additionally received a percutaneous drainage. Three patients were forwarded to surgery due to necrosis: one necrosectomy (idiopathic pancreatitis), one gastrojejunostomy (alcoholic pancreatitis) and one removal of dislocated transgastral drainage (alcoholic pancreatitis). Hence, an intervention was performed in a total of eleven episodes (2.8%).

The distribution of age and gender among the four etiology groups is shown in Figure 1. In biliary pancreatitis incidence increased with age. The group was dominated by women and the incidence peaked in the 4th and 6th decade. In alcoholic pancreatitis there was a marked peak in the 5th decade and this group was dominated by men. In idiopathic pancreatitis there was a peak in the 8th and possibly in the 5th decade, the men were younger than the women. In the group with miscellaneous causes there was no identifiable pattern.

Other characteristics of the different etiologies are given in Table 1. There were highly significant differences between the etiologies with regard to the baseline parameters gender, age, percentage of recurrent episodes, evidence of chronic pancreatitis, and plasma lipase on admission. There were also significant differences with regard to body mass, heart rate, and hematocrit. No differences could be detected with regard to comorbidity, systolic blood pressure, plasma creatinine, and white blood cell count. The outcome parameters including persistent organ




Figure 1. NAPOLI-1 Study Schema

failure, occurrence of necrosis, and mortality differed significantly, maximum C-reactive protein (CRP) and duration of hospitalization did not.

Recurrent episodes and chronic pancreatitis were frequent in alcoholic pancreatitis. When relapses were excluded the lipase in alcoholic pancreatitis was still comparably low ($1,540 \pm 241$ U/L; $n=61$). The same was true when patients with chronic pancreatitis were excluded in this group ($1,297 \pm 191$ U/L; $n=72$).

DISCUSSION

To our knowledge this is the first study on a large number of consecutive episodes of acute pancreatitis with a thorough work-up of etiology in all cases. Previous studies have defined biliary pancreatitis by non-specific methods [8, 11, 12, 14] or did not give any definition [7, 9]. We therefore assume that biliary origin might not have been categorized correctly in all cases. This would in part explain the discrepant conclusions on characteristics of the etiological groups. In our study, we used a number of tests to define the correct entity [17, 21]. Moreover, the vast majority of patients underwent EUS to exclude undetected choledocholithiasis and to determine abnormalities of the pancreas.

Our data shows that different etiologies of acute pancreatitis differ in a variety of aspects. Firstly, the characteristics of the four groups differ significantly. As shown in Figure 1 we were able to reproduce the well-established features of age and gender in biliary and alcoholic pancreatitis [6, 7]. The marked differences in body mass between these two groups may be explained by the association of gallstones with obesity and the high prevalence of chronic pancreatitis (and probably also malnutrition) in patients with alcoholic episodes. The characteristics of idiopathic pancreatitis have been specified less clearly. In a large epidemiological study, patients with idiopathic pancreatitis were of same age and similar gender distribution as patients with pancreatitis of biliary origin [7]. However, the authors point out the possibility of misclassifications by overlooking gallstones and, hence, misinterpreting biliary pancreatitis as idiopathic. In our study we found a frequency peak in the 8th decade and possibly also in the 5th. This pattern contrasts with the cases of non-biliary, non-alcoholic pancreatitis caused by an obvious trigger where gender and age were distributed evenly.

Our data also indicate that alcohol is the leading cause for recurrent episodes followed by idiopathic etiology and other specific causes. This is consistent with observations of the natural course of acute pancreatitis showing idiopathic etiology, pancreas divisum, cigarette smoking and alcohol consumption to be associated with relapse [22-24]. The rate of relapses in biliary pancreatitis was rather small probably due to a timely cholecystectomy in most cases. The rate of chronic pancreatitis was considerable in idiopathic pancreatitis contradicting the notion that progression from acute to chronic pancreatitis only occurs in alcoholics [4]. Alternatively, these episodes could have been flare-ups of idiopathic chronic pancreatitis.

The second aspect is the presentation on admission. There were significant differences in heart rate, plasma lipase, and haematocrit. Biliary pancreatitis was associated with a higher, alcoholic pancreatitis with a lower lipase. This discrepancy has already been observed in other studies [25-26]. The difference was still evident when we excluded patients with recurrent or chronic pancreatitis. It therefore seems unlikely that the lower lipase in alcoholic pancreatitis is attributable to a loss of exocrine capacity over a longer period of organ damage. It can rather be explained by a different dynamic of a toxic trigger in contrast to the sudden incident of gallstone obstruction. It might also be attributed to the analgesic effect of alcohol leading to a delay in presentation to the emergency department [27].

Patients with alcoholic pancreatitis had the highest heart rate and the highest hematocrit on admission. Both parameters are indicators for hemoconcentration. Hemoconcentration on admission has been linked to the development of subsequent necrosis [28, 29]. In our study, patients with an alcoholic episode indeed had the highest rate of necrosis. In contrast, the necrosis rate in

Table 1. Characteristics of the episodes of acute pancreatitis sorted by etiology. P values for differences among the groups. The continuous variables are given as mean \pm standard deviation.

	Biliary (n=)	Alcoholic (n=)	Idiopathic (n=)	Others (n=)	P value^a
Gender (male/female)	57/88	98/25	29/32	29/33	<0.001
Age (years)	66.5 \pm 18.7	50.2 \pm 12.2	61.1 \pm 17.4	52.2 \pm 17.7	<0.001
Body mass index (kg/m ²)	28.7 \pm 10.7	25.2 \pm 5.4	27.9 \pm 4.9	26.3 \pm 5.9	0.004
CCI \geq 3	24.8%	18.7%	24.6%	32.3%	0.233
Recurrent episode (%)	11.7%	50.4%	36.1%	33.6%	<0.001
Chronic pancreatitis (%)	0.7%	41.5%	9.8%	16.1%	<0.001
Heart rate (min ⁻¹)	79 \pm 19	87 \pm 20	78 \pm 21	84 \pm 19	0.004
Systolic blood pressure (mmHg)	138 \pm 25	142 \pm 25	138 \pm 22	134 \pm 23	0.144
Lipase (U/L)	4,591 \pm 5,131	1,186 \pm 1,553	2,448 \pm 3,612	1,761 \pm 2,616	<0.001
Hematocrit (%)	39 \pm 4	41 \pm 5	39 \pm 6	38 \pm 6	0.009
Plasma creatinine (mmol/L)	88 \pm 68	90 \pm 106	88 \pm 84	79 \pm 51	0.087
White blood cell count (nl ⁻¹)	11.9 \pm 5.0	12.6 \pm 4.6	12.9 \pm 5.0	11.4 \pm 5.3	0.128
Persistent organ failure (%)	2.8%	4.1%	16.4%	3.2%	0.005
Necrosis (%)	8.3%	20.3%	16.4%	11.3%	0.032
Maximum CRP (mg/L)	105 \pm 92	125 \pm 106	126 \pm 109	116 \pm 130	0.409
Hospitalization (days)	8.1 \pm 5.7	10.1 \pm 10.4	10.8 \pm 12.3	10.2 \pm 7.0	0.436
Mortality	1.4%	0.8%	9.8%	4.8%	0.004

CCI: Charlson comorbidity index; CRP: C-reactive protein

^aSignificant P values are shown in bold

idiopathic pancreatitis was nearly as high although heart rate and hematocrit on admission were as low as those in pancreatitis of biliary origin. Hence, the association might be an epiphenomenon combining two distinct features of alcoholic pancreatitis. This could offer an explanation for the finding, that aggressive volume therapy does not prevent necrosis [20, 30].

The third aspect is the outcome. While alcoholic pancreatitis had the highest rate of necrosis and a low mortality, idiopathic pancreatitis had by far the highest rates of persistent organ failure and also the highest mortality. This is in line with the observations that necrosis and organ failure are independent determinants and that persistent organ failure indicates a poor outcome [31, 32]. The high mortality in idiopathic acute pancreatitis has also been observed in a number of previous studies [7, 14-16]. We therefore conclude that identifying a cause for acute pancreatitis could be helpful to exclude patients at high risk for systemic complications.

CONCLUSION

In summary, we found substantial differences between the characteristics of the major etiological types of acute pancreatitis. Alcoholic episodes have the highest risk of necrosis and the worst outcome can be found in the idiopathic group. Differentiating between the etiologies seems to be necessary to correctly assess the risk for certain complications. Large scaled studies with a thorough search for the cause in each episode will be necessary to determine specific risk factors.

Conflict of Interest

Authors declare to have no conflict of interest.

References

- Wang GJ, Gao CF, Wei D, Wang C, Ding SQ. Acute pancreatitis: etiology and common pathogenesis. *World J Gastroenterol* 2009; 15:1427-30. [PMID: 19322914]
- Kylänpää L, Rakonczay Z, Jr., O'Reilly DA. The clinical course of acute pancreatitis and the inflammatory mediators that drive it. *Int J Inflam* 2012; 2012:360685. [PMID: 23304633]
- Gullo L, Migliori M, Pezzilli R, Olah A, Farkas G, Levy P et al. An update on recurrent acute pancreatitis: data from five European countries. *Am J Gastroenterol* 2002; 97:1959-62. [PMID: 12190160]
- Lankisch PG, Breuer N, Bruns A, Weber-Dany B, Löwenfels AB, Maisonneuve P. Natural history of acute pancreatitis: a long-term population-based study. *Am J Gastroenterol* 2009; 104:2797-805. [PMID: 19603011]
- Zheng Y, Zhou Z, Li H, Li J, Li A, Ma B et al. A Multicenter Study on Etiology of Acute Pancreatitis in Beijing During 5 Years. *Pancreas* 2014. [PMID: 25438072]
- Lankisch PG, Assmus C, Maisonneuve P, Löwenfels AB. Epidemiology of pancreatic diseases in Lüneburg County. A study in a defined german population. *Pancreatol* 2002; 2:469-77. [PMID: 12378115]
- Frey CF, Zhou H, Harvey DJ, White RH. The incidence and case-fatality rates of acute biliary, alcoholic, and idiopathic pancreatitis in California, 1994-2001. *Pancreas* 2006; 33:336-44. [PMID: 17079936]
- Uhl W, Isenmann R, Curti G, Vogel R, Beger HG, Buchler MW. Influence of etiology on the course and outcome of acute pancreatitis. *Pancreas* 1996; 13:335-43. [PMID: 8899793]
- Uomo G, Pezzilli R, Gabbriellini A, Castoldi L, Zerbi A, Frulloni L et al. Diagnostic assessment and outcome of acute pancreatitis in Italy: results of a prospective multicentre study. *ProInf-AISP: Progetto informatizzato pancreatite acuta, Associazione Italiana Studio Pancreas, phase II. Dig Liver Dis* 2007; 39:829-37. [PMID: 17625994]
- Choi JH, Kim MH, Oh D, Paik WH, Park dH, Lee SS et al. Clinical relevance of the revised Atlanta classification focusing on severity stratification system. *Pancreatol* 2014; 14:324-9. [PMID: 25174301]
- Omdal T, Dale J, Lie SA, Iversen KB, Flaatten H, Ovrebø K. Time trends in incidence, etiology, and case fatality rate of the first attack of acute pancreatitis. *Scand J Gastroenterol* 2011; 46:1389-98. [PMID: 21830851]
- Lankisch PG, Assmus C, Pflüthofer D, Struckmann K, Lehnick D. Which etiology causes the most severe acute pancreatitis? *Int J Pancreatol* 1999; 26:55-7. [PMID: 10597400]

13. Talukdar R, Clemens M, Vege SS. Moderately severe acute pancreatitis: prospective validation of this new subgroup of acute pancreatitis. *Pancreas* 2012; 41:306-9. [PMID: 22015971]
14. de Beaux AC, Palmer KR, Carter DC. Factors influencing morbidity and mortality in acute pancreatitis; an analysis of 279 cases. *Gut* 1995; 37:121-6. [PMID: 7672660]
15. Bai Y, Liu Y, Jia L, Jiang H, Ji M, Lv N et al. Severe acute pancreatitis in China: etiology and mortality in 1976 patients. *Pancreas* 2007; 35:232-7. [PMID: 17895843]
16. Chen Y, Zak Y, Hernandez-Boussard T, Park W, Visser BC. The epidemiology of idiopathic acute pancreatitis, analysis of the nationwide inpatient sample from 1998 to 2007. *Pancreas* 2013; 42:1-5. [PMID: 22750972]
17. Ortega AR, Gomez-Rodriguez R, Romero M, Fernandez-Zapardiel S, Cespedes MM, Carrobes JM. Prospective comparison of endoscopic ultrasonography and magnetic resonance cholangiopancreatography in the etiological diagnosis of "idiopathic" acute pancreatitis. *Pancreas* 2011; 40:289-94. [PMID: 21206330]
18. Freeman ML, Werner J, van Santvoort HC, Baron TH, Besselink MG, Windsor JA et al. Interventions for necrotizing pancreatitis: summary of a multidisciplinary consensus conference. *Pancreas* 2012; 41:1176-94. [PMID: 23086243]
19. Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG et al. Classification of acute pancreatitis - 2012: revision of the Atlanta classification and definitions by international consensus. *Gut* 2013; 62:102-11. [PMID: 23100216]
20. Weitz G, Woitalla J, Wellhoner P, Schmidt K, Buning J, Fellermann K. Detrimental effect of high volume fluid administration in acute pancreatitis - A retrospective analysis of 391 patients. *Pancreatol* 2014; 14(6):478-483.
21. Perez-Mateo M. How we predict the etiology of acute pancreatitis. *JOP* 2006; 7:257-61. [PMID: 25451185]
22. Yadav D, O'Connell M, Papachristou GI. Natural history following the first attack of acute pancreatitis. *Am J Gastroenterol* 2012; 107:1096-103. [PMID: 22613906]
23. Vippera K, Papachristou GI, Easler J, Muddana V, Slivka A, Whitcomb DC et al. Risk of and factors associated with readmission after a sentinel attack of acute pancreatitis. *Clin Gastroenterol Hepatol* 2014; 12:1911-9. [PMID: 24815327]
24. Cavestro GM, Leandro G, Di Leo M, Zupparado RA, Morrow OB, Notaristefano C et al. A single-centre prospective, cohort study of the natural history of acute pancreatitis. *Dig Liver Dis* 2014. [PMID: 25475611]
25. Pezzilli R, Billi P, Miglioli M, Gullo L. Serum amylase and lipase concentrations and lipase/amylase ratio in assessment of etiology and severity of acute pancreatitis. *Dig Dis Sci* 1993; 38:1265-9. [PMID: 7686843]
26. Cornett DD, Spier BJ, Eggert AA, Pfau PR. The causes and outcome of acute pancreatitis associated with serum lipase >10,000 u/l. *Dig Dis Sci* 2011; 56:3376-81. [PMID: 21614591]
27. James MF, Duthie AM, Duffy BL, McKeag AM, Rice CP. Analgesic effect of ethyl alcohol. *Br J Anaesth* 1978; 50:139-41. [PMID: 341934]
28. Baillargeon JD, Orav J, Ramagopal V, Tenner SM, Banks PA. Hemoconcentration as an early risk factor for necrotizing pancreatitis. *Am J Gastroenterol* 1998; 93:2130-4. [PMID: 9820385]
29. Gan SI, Romagnuolo J. Admission hematocrit: a simple, useful and early predictor of severe pancreatitis. *Dig Dis Sci* 2004; 49:1946-52. [PMID: 15628731]
30. de Madaria E, Soler-Sala G, Sanchez-Paya J, Lopez-Font I, Martinez J, Gomez-Escolar L et al. Influence of fluid therapy on the prognosis of acute pancreatitis: a prospective cohort study. *Am J Gastroenterol* 2011; 106:1843-50. [PMID: 21876561]
31. Lankisch PG, Pflithofer D, Lehnick D. No strict correlation between necrosis and organ failure in acute pancreatitis. *Pancreas* 2000; 20:319-22. [PMID: 10766460]
32. Thandassery RB, Yadav TD, Dutta U, Appasani S, Singh K, Kochhar R. Dynamic nature of organ failure in severe acute pancreatitis: the impact of persistent and deteriorating organ failure. *HPB (Oxford)* 2013; 15:523-8. [PMID: 23750495]